

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
25 September 2003 (25.09.2003) PCT
(10) International Publication Number
WO 03/078445 A2(51) International Patent Classification: C07H 21/00
(21) International Application Number: PCT/DK03/00173
(22) International Filing Date: 14 March 2003 (14.03.2003)(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
PA 2002 0415 15 March 2002 (15.03.2002) DK
60364.056 15 March 2002 (15.03.2002) US
PA 2002 01952 19 December 2002 (19.12.2002) DK
60434.429 19 December 2002 (19.12.2002) US(71) Applicant (for all designated States except US): NUEVO-
LUTION A/S [DK/DK]; Rønnegade 8, 5th floor, DK-2100
Copenhagen Ø (DK).(72) Inventors; and
(75) Inventors/Applicants (for US only): GOULIAEV, Alex,
Hahr (DK/DK); Brandsted 223, DK-3670 Veksø Sjael-
land (DK); PEDERSEN, Henrik (DK/DK); Prodesvej 24,
DK-2880 Bagsværd (DK); GODSKESSEN, Michael, Au-
ders (DK/DK); Plantagekrogen 8, DK-2950 Vedbaek (DK);
KRONBORG, Tue, Titthol, Alstenleim (DK/DK);
Ndr. Frihavnsgade 94, 1.v., DK-2100 København Ø (DK);
HUSEMOEN, Gitte, Nystrup (DK/DK); Bregnerødsgade
18, 1.th., DK-2200 København N (DK); HO, JustinPublished:
— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: A BUILDING BLOCK FORMING A C=C DOUBLE BOND UPON REACTION

(57) Abstract: A building block having the dual capabilities of recognising an encoding element and transferring a Functional Entity
Precursor to a recipient reactive group is disclosed. The building block may be used in the generation of a single complex or libraries
of different complexes, wherein the complex comprises an encoded molecule linked to an encoding element. Libraries of complexes
are useful in the quest for pharmaceutically active compounds.

Title

A BUILDING BLOCK FORMING A C=C DOUBLE BOND UPON REACTION

Technical Field of the Invention

5 The present invention relates to a building block comprising a complementing ele-
ment and a precursor for a functional entity. The building block is designed to trans-
fer the Functional Entity Precursor to a recipient reactive group upon recognition
between the complementing element and an encoding element associated with the
reactive group.

Background

10 The transfer of a chemical entity from one mono-, di- or oligonucleotide to another
has been considered in the prior art. Thus, N. M. Chung *et al.* (Biochim. Biophys.
Acta, 1971, 228, 536-543) used a poly(U) template to catalyse the transfer of an ace-
tyl group from 3'-O-acetyladenosine to the 5'-OH of adenosine. The reverse transfer,
i.e. the transfer of the acetyl group from a 5'-O-acetyladenosine to a 3'-OH group of
another adenosine, was also demonstrated.

15 20 25
Walder *et al.* Proc. Natl. Acad. Sci. USA, 1979, 76, 51-55 suggest a synthetic pro-
cedure for peptide synthesis. The synthesis involves the transfer of nascent immobi-
lized polypeptide attached to an oligonucleotide strand to a precursor amino acid
attached to an oligonucleotide. The transfer comprises the chemical attack of the
amino group of the amino acid precursor on the substitution labile peptidyl ester,
which in turn results in an acyl transfer. It is suggested to attach the amino acid pre-
cursor to the 5' end of an oligonucleotide with a thiol ester linkage.

30 The transfer of a peptide from one oligonucleotide to another using a template is
disclosed in Bruick RK *et al.* Chemistry & Biology, 1996, 3:49-56. The carboxy ter-
minal of the peptide is initially converted to a thioester group and subsequently
transformed to an activated thioester upon incubation with Ellman's reagent. The
activated thioester is reacted with a first oligo, which is 5'-thiol-terminated, resulting
in the formation of a thio-ester linked intermediate. The first oligonucleotide and a
second oligonucleotide having a 3' amino group is aligned on a template such that
the thioester group and the amino group are positioned in close proximity and a

transfer is effected resulting in a coupling of the peptide to the second oligonucleotide through an amide bond.

- 5 The Wittig reaction (Wittig and Geissler, Liebigs Ann. Chem. (1953), 580, 44) is a well appreciated C=C double bond forming reaction in organic synthesis (Chem. Rev. (1989), 89, 863). The reaction with stabilised phosphonium salts is however limited to reaction with primarily aldehydes and furthermore favour formation of only Z-alkenes. The Horner-Wadsworth-Emmons (HWE) modification of the Wittig reaction provides several advantages. Firstly, the reactive species of stabilised HWE-reagents is more reactive and therefore reads with a broader range of carbonyl compounds, even hindered ketones (Comprehensive Organic Synthesis, Vol I, p. 762). Secondly, the stereochemical identity of the formed C=C double bond can be controlled by the substituents on the phosphorous atom. Formation of Z-alkenes are favoured by using Stills or Andos HWE-reagents (Still and Gennari, Tetrahedron Lett. (1983), 24, 4405-08, Ando, J. Org. Chem. (1998), 63, 8411-8416), whereas the formation of E-alkenes are favoured using simple alkyl based HWE-reagents (Chem. Rev. (1989), 89, 863).

Summary of the invention


20 The present invention relates to a building block of the general formula:

Complementing Element - Linker - Carrier - Functional entity precursor capable of transferring a functional entity precursor to a recipient reactive group, wherein

Complementing Element is a group identifying the functional entity,

Linker is a chemical moiety comprising a spacer and a S-C-connecting group, wherein the spacer is a valence bond or a group distancing the functional entity precursor to be transferred from the complementing element and the S-C-connecting group connects the spacer with the Carrier,

Carrier is selected among the groups consisting of



wherein the Functional entity precursor is attached to the phosphorous atom,

each individual X is chosen independently among the group consisting of NR¹⁶, O, S, and Se;

U = R³,

5 R¹ = H, C₁-C₈ alkyl, C₁-C₈ hydroxyalkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, aryl or heteroaryl; optionally substituted with one or more F, Cl, Br, I, or CN, and where each of the individual R¹ groups are chosen independently;

R² = a valence bond, C₁-C₈ Alkylene, C₂-C₈ Alkenylene, C₂-C₈ Alkynylene, Arylene, Heteroarylene, C₁-C₈ Alkylene-arylene or C₁-C₈ Alkylene-heteroarylene;

10 R³ is selected from -H, -OR⁴, -NR⁴, -F, -Cl, -Br, -I, -NO₂, -CN, -C(Halogen)₃, -C(O)R⁴, -C(O)NHR⁴, C(O)NR⁴, -NC(O)R⁴, -S(O)₂NHR⁴, -S(O)₂NR⁴, -S(O)₂R⁴, -P(O)₂R⁴, -P(O)₂R⁴, -S(O)₂R⁴, P(O)-OR⁴, -S(O)-OR⁴, -N⁺R⁴, wherein and each individual R⁴ is chosen independently from H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, or aryl,

R¹⁶ is selected independently from H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, aryl, C₁-C₈ al-

15 kylene-aryl, $\begin{matrix} \text{G} \\ | \\ \text{---O---} \\ | \\ \text{---G} \end{matrix}$ or $\begin{matrix} \text{G} \\ | \\ \text{---N---} \\ | \\ \text{---G} \end{matrix}$, G is H or C₁-C₈ alkyl and n is 1, 2, 3 or 4.

Functional entity precursor is of the general formula C(H)(V)-W;

20 wherein V independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloalkadienyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of SnR⁵R⁶R⁷, Sn(OR⁵)R⁶R⁷, Sn(OR⁵)(OR⁶)R⁷, BR⁵R⁶, B(OR⁵)R⁶, B(OR⁵)(OR⁶), halogen, CN, CNO, C(halogen)₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, N₃, NR⁵R⁶, N⁺R⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, NC, P(=O)(OR⁵)OR⁶, P⁺R⁵R⁶R⁷, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶, C(=O)NR⁵R⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶, wherein,

25 R⁵, R⁶ and R⁷ independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloalkadienyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CNO, C(halogen)₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, N₃, NR⁵R⁶, N⁺R⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, NC, P(=O)(OR⁵)OR⁶, P⁺R⁵R⁶R⁷, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)OR⁵, C(=O)NR⁵R⁶,

$C(=O)NR^8OR^9C(=NR^5)NR^6R^7$, $C(=O)NR^5NR^6R^7$ or $C(=O)NR^5NR^6R^{10}$, wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring.

5

wherein,
 R^8 , R^9 and R^{10} independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl and wherein R^8 and R^9 may together form a 3-8 membered heterocyclic ring or R^6 and R^{10} may together form a 3-8 membered heterocyclic ring or R^6 and R^{10} may together form a 3-8 membered heterocyclic ring.

10

W is selected among the group consisting of H, aryl, heteroaryl, $C(=O)OR^{11}$, $C(O)R^{11}$, $C(=O)NR^{11}$, $C(=O)NR^{11}OR^{11}$, $C(=NR^{11})OR^{11}$, $C(=NOR^{11})R^{11}$, $C(=NNR^{11})R^{11}$, $S(O)R^{11}$, $S(O)_2R^{11}$, $S(O)NR^{11}$, $S(O)_2NR^{11}$, -CN, $P(O)R^{11}$, -NO₂, NR^{11} , R^3 or SR^{11} .

15 where each individual R^{11} is H or selected independently among the group consisting of C_1 -C₆ alkyl, C_2 -C₆ alkenyl, C_2 -C₆ alkynyl, C_3 -C₇ cycloalkyl, C_3 -C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R^{12} and 0-3 R^{15} and each individual R^{12} is selected independently from -N₃, -CNO, -C(NOH)NH₂, -NHOH, -NHR⁴NHR⁴, -C(O)R⁴, -SnR⁴, -B(OR⁴)₂, -P(O)(OR⁴)₂ or the group consisting of C_2 -C₆ alkynyl, C_2 -C₆ alkynyl, C_1 -C₆ alkadienyl said group being substituted with 0-2 R^{13} .

20

where each individual R^{13} is independently selected from -NO₂, -C(O)OR¹⁴, -C(O)R¹⁴, -CN, -OSiR¹⁴, -OR¹⁴ and -NR¹⁴.

25 each individual R^{14} is independently chosen from a group comprising H, C_1 -C₆ alkyl, C_2 -C₆ alkenyl, C_2 -C₆ alkynyl, C_3 -C₇ cycloalkyl, aryl or C_1 -C₆ alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -NO₂, -R⁴, -OR⁴, -SiR⁴.

R^{15} is -O, -F, -Cl, -Br, -I, -CN, -NO₂, -OR¹⁴, -NR¹⁴, -NR¹⁴-C(O)R¹⁴, -NR¹⁴-C(O)OR¹⁴, -SR¹⁴, -S(O)R¹⁴, -S(O)₂R¹⁴, -COOR¹⁴, -C(O)NR¹⁴, and -S(O)₂NR¹⁴.

30

The term " C_3 -C₇ cycloheteroalkyl" as used herein refers to a radical of totally saturated heterocycle like a cyclic hydrocarbon containing one or more heteroatoms selected from nitrogen, oxygen, phosphorus, boron and sulphur independently in the cycle such as pyrrolidine (1-pyrrolidine; 2-pyrrolidine; 3-pyrrolidine; 4-pyrrolidine; 5-pyrrolidine); pyrazolidine (1-pyrazolidine; 2-pyrazolidine; 3-pyrazolidine; 4-pyrazolidine; 5-pyrazolidine); imidazolidine (1-imidazolidine; 2-imidazolidine; 3-imidazolidine; 4-imidazolidine; 5-imidazolidine); thiazolidine (2-thiazolidine; 3-thiazolidine; 4-thiazolidine; 5-thiazolidine); piperidine (1-piperidine; 2-piperidine; 3-piperidine; 4-piperidine; 5-piperidine; 6-piperidine); piperazine (1-piperazine; 2-piperazine; 3-piperazine; 4-piperazine; 5-piperazine; 6-piperazine); morpholine (2-morpholine; 3-morpholine; 4-morpholine; 5-morpholine; 6-morpholine); thiomorpholine (2-thiomorpholine; 3-thiomorpholine; 4-thiomorpholine; 5-thiomorpholine; 6-thiomorpholine); 1,2-oxathiolane (3-(1,2-oxathiolane); 4-(1,2-oxathiolane); 5-(1,2-oxathiolane); 1,3-dioxolane (2-(1,3-dioxolane); 4-(1,3-dioxolane); 5-(1,3-dioxolane); tetrahydropyran; (2-tetrahydropyran; 3-tetrahydropyran; 4-tetrahydropyran; 5-tetrahydropyran; 6-tetrahydropyran); hexahydropyridazine (1-hexahydropyridazine); 2-(hexahydropyridazine); 3-(hexahydropyridazine); 4-(hexahydropyridazine); 5-(hexahydropyridazine); 6-(hexahydropyridazine)), [1,3,2]dioxaborolane, [1,3,6,2]dioxazaborocane

15

The term "aryl" as used herein includes carbocyclic aromatic ring systems of 5-7 carbon atoms. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems as well as up to four fused fused aromatic- or partially hydrogenated rings, each ring comprising 5-7 carbon atoms.

20

The term "heteroaryl" as used herein includes heterocyclic unsaturated ring systems containing, in addition to 2-18 carbon atoms, one or more heteroatoms selected from nitrogen, oxygen and sulphur such as furyl, thienyl, pyrrolyl, heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated below.

25

The terms "aryl" and "heteroaryl" as used herein refers to an aryl which can be optionally substituted or a heteroaryl which can be optionally substituted and includes phenyl, biphenyl, indenyl, naphthyl (1-naphthyl, 2-naphthyl), N-hydroxytetrazolyl, N-hydroxytriazolyl, N-hydroxyimidazolyl, anthracenyl (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), thiophenyl (2-thienyl, 3-thienyl), furyl (2-furyl, 3-furyl), indolyl, oxadiazolyl, isoxazolyl, quinazolyl, fluorenyl, xanthenyl, isindanylyl, benzhydryl, acridinyl, thiazolyl, pyrrolyl (2-pyrrolyl), pyrazolyl (3-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (2-pyrimidinyl, 4-

30

35

zolidine; 4-pyrazolidine; 5-pyrazolidine); imidazolidine (1-imidazolidine; 2-imidazolidine; 3-imidazolidine; 4-imidazolidine; 5-imidazolidine); thiazolidine (2-thiazolidine; 3-thiazolidine; 4-thiazolidine; 5-thiazolidine); piperidine (1-piperidine; 2-piperidine; 3-piperidine; 4-piperidine; 5-piperidine; 6-piperidine); piperazine (1-piperazine; 2-piperazine; 3-piperazine; 4-piperazine; 5-piperazine; 6-piperazine); morpholine (2-morpholine; 3-morpholine; 4-morpholine; 5-morpholine; 6-morpholine); thiomorpholine (2-thiomorpholine; 3-thiomorpholine; 4-thiomorpholine; 5-thiomorpholine; 6-thiomorpholine); 1,2-oxathiolane (3-(1,2-oxathiolane); 4-(1,2-oxathiolane); 5-(1,2-oxathiolane); 1,3-dioxolane (2-(1,3-dioxolane); 4-(1,3-dioxolane); 5-(1,3-dioxolane); tetrahydropyran; (2-tetrahydropyran; 3-tetrahydropyran; 4-tetrahydropyran; 5-tetrahydropyran; 6-tetrahydropyran); hexahydropyridazine (1-hexahydropyridazine); 2-(hexahydropyridazine); 3-(hexahydropyridazine); 4-(hexahydropyridazine); 5-(hexahydropyridazine); 6-(hexahydropyridazine)), [1,3,2]dioxaborolane, [1,3,6,2]dioxazaborocane

The term "aryl" as used herein includes carbocyclic aromatic ring systems of 5-7 carbon atoms. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems as well as up to four fused fused aromatic- or partially hydrogenated rings, each ring comprising 5-7 carbon atoms.

The term "heteroaryl" as used herein includes heterocyclic unsaturated ring systems containing, in addition to 2-18 carbon atoms, one or more heteroatoms selected from nitrogen, oxygen and sulphur such as furyl, thienyl, pyrrolyl, heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated below.

The terms "aryl" and "heteroaryl" as used herein refers to an aryl which can be optionally substituted or a heteroaryl which can be optionally substituted and includes phenyl, biphenyl, indenyl, naphthyl (1-naphthyl, 2-naphthyl), N-hydroxytetrazolyl, N-hydroxytriazolyl, N-hydroxyimidazolyl, anthracenyl (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), thiophenyl (2-thienyl, 3-thienyl), furyl (2-furyl, 3-furyl), indolyl, oxadiazolyl, isoxazolyl, quinazolyl, fluorenyl, xanthenyl, isindanylyl, benzhydryl, acridinyl, thiazolyl, pyrrolyl (2-pyrrolyl), pyrazolyl (3-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (2-pyrimidinyl, 4-

pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl), pyrazinyl, pyridazinyl (3- pyridazinyl, 4- pyridazinyl, 5-pyridazinyl), quinolyl (2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6- quinolyl, 7-quinolyl, 8-quinolyl), isquinolyl (1-isquinolyl, 3-isquinolyl, 4- isquinolyl, 5-isquinolyl, 6-isquinolyl, 7-isquinolyl, 8-isquinolyl), benzo[b]furanyl (2-benzo[b]furanyl, 3-benzo[b]furanyl, 4-benzo[b]furanyl, 5- benzo[b]furanyl, 6-benzo[b]furanyl, 7-benzo[b]furanyl), 2,3-dihydro-benzo[b]furanyl (2-(2,3-dihydro-benzo[b]furanyl), 3-(2,3-dihydro-benzo[b]furanyl), 4-(2,3-dihydro-benzo[b]furanyl), 5-(2,3-dihydro-benzo[b]furanyl), 6-(2,3-dihydro-benzo[b]furanyl), 7-(2,3-dihydro-benzo[b]furanyl), benzo[b]thiophenyl (2-benzo[b]thiophenyl, 3- benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6- benzo[b]thiophenyl, 7-benzo[b]thiophenyl), 2,3-dihydro-benzo[b]thiophenyl (2-(2,3- dihydro-benzo[b]thiophenyl), 3-(2,3-dihydro-benzo[b]thiophenyl), 4-(2,3-dihydro- benzo[b]thiophenyl), 5-(2,3-dihydro-benzo[b]thiophenyl), 6-(2,3-dihydro- benzo[b]thiophenyl), 7-(2,3-dihydro-benzo[b]thiophenyl), 8-(2,3-dihydro- benzo[b]thiophenyl), 9-(2,3-dihydro-benzo[b]thiophenyl), indolyl (1-indolyl, 2- indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), indazole (1-indazolyl, 3- indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl), benzimidazolyl (1- benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl, 5-benzimidazolyl, 6- benzimidazolyl, 7-benzimidazolyl, 8-benzimidazolyl), benzoxazolyl (1- benzoxazolyl, 2-benzoxazolyl), benzothiazolyl (1-benzothiazolyl, 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl), carbazolyl (1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl), 5H-dibenz[b, f]azepine (5H- dibenz[b, f]azepine-1-yl, 5H-dibenz[b, f]azepine-2-yl, 5H-dibenz[b, f]azepine-3-yl, 5H- dibenz[b, f]azepine-4-yl, 5H-dibenz[b, f]azepine-5-yl), 10,11-dihydro-5H- dibenz[b, f]azepine (10,11-dihydro-5H-dibenz[b, f]azepine-1-yl, 10,11-dihydro-5H- dibenz[b, f]azepine-2-yl, 10,11-dihydro-5H-dibenz[b, f]azepine-3-yl, 10,11-dihydro- 5H-dibenz[b, f]azepine-4-yl, 10,11-dihydro-5H-dibenz[b, f]azepine-5-yl).

The term "halogen" designates an atom selected from the group consisting of -F, - Cl, -Br and -I.

30 In the following description of the invention the direction of connections between the various components of a building block should be read left to right. For example an S-C-connecting group -C(=O)-NH- is connected to a Spacer through the carbon atom on the left and to a Carrier through the nitrogen atom on the right hand side.

5 The Functional Entity carries elements used to interact with host molecules and optionally reactive elements allowing further elaboration of an encoded molecule of a library. Interaction with host molecules like enzymes, receptors and polymers is typically mediated through van der Waals' interactions, polar- and ionic interactions and pi-stacking effects. Substituents mediating said effects may be masked by methods known to an individual skilled in the art (Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*; 3rd ed.; John Wiley & Sons: New York, 1999) to avoid undesired interactions or reactions during the preparation of the individual building blocks and during library synthesis. Analogously, reactive elements may be masked by suitably selected protection groups. It is appreciated by one skilled in the art that by suitable protection, a functional entity may carry a wide range of substituents.

10 The Functional Entity Precursor may be a masked Functional Entity that is incorporated into an encoded molecule. After incorporation, reactive elements of the Functional Entity may be revealed by un-masking allowing further synthetic operations. Finally, elements mediating recognition of host molecules may be un-masked.

20 In a certain aspect of the invention the Functional entity precursor is of the general formula C(H)(V)-W; in which V is H, -F, -Cl, -Br, -I, -CN, -NO₂ or -OR¹⁴, or selected among the group consisting of C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, -O-aryl and -O-heteroaryl said group being substituted with 0-3 R¹¹, 0-3 R¹² and 0-3 R¹⁵, or V is C₁-C₃ alkylene-NR¹¹, C₁-C₃ alkylene-NR¹¹C(O)R¹⁴, C₁-C₃ alkylene-NR¹¹C(O)OR¹⁴, C₁-C₃ alkylene-O-NR¹¹, C₁-C₃ alkylene-O-NR¹¹C(O)R¹⁴, C₁-C₃ alkylene-O-NR¹¹C(O)OR¹⁴ substituted with 0-3 R¹⁵, W is selected among the group consisting of H, aryl, heteroaryl, C(=O)OR¹¹, C(O)R¹¹, C(=O)NR¹¹, C(=O)NR¹¹OR¹¹, C(=NR¹¹)OR¹¹, C(=NOR¹¹)R¹¹, C(=NNR¹¹)₂R¹¹, S(O)R¹¹, S(O)₂R¹¹, S(O)NR¹¹, S(O)₂NR¹¹, -CN, P(O)R¹¹, -NO₂, NR¹¹,⁺ or SR¹¹,⁺



30 Where each individual R¹¹ is H or selected independently among the group consisting of C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R¹² and 0-3 R¹⁵ and Each individual R¹² is selected independently from -N₃, -CNO, -C(NOH)NH₂, -NHOH, -NHR¹⁴NHR¹⁴, -C(O)R¹⁴, -SnR¹⁴, -B(OR¹⁴)₂, -P(O)(OR¹⁴)₂ or the group consist-

ing of C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₆ alkadienyl said group being substituted with 0-2 R¹³,

where R¹³ is independently selected from -NO₂, -C(O)OR¹⁴, -C(O)R¹⁴, -CN, -OSiR¹⁴₃, -OR¹⁴ and -NR¹⁴.

5 Each individual R¹⁴ is independently chosen from a group comprising H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -NO₂, -R⁴, -OR⁴, -SiR⁴

10 R¹⁵ is -O-, -F-, -Cl-, -Br-, -I-, -CN, -NO₂, -OR¹⁴, -NR¹⁴₂, -NR¹⁴₂-C(O)R¹⁴, -NR¹⁴₂-C(O)OR¹⁴, -SR¹⁴, -S(O)R¹⁴, -S(O)₂R¹⁴, -COOR¹⁴, -C(O)NR¹⁴₂ and -S(O)₂NR¹⁴₂.

R¹⁶ is selected independently from H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl, C₁-C₆ alkylene-aryl,  or , G is H or C₁-C₆ alkyl and n is 1, 2, 3 or 4.


15 In a preferred embodiment of the present invention, W is selected among the group consisting of C(=O)OR¹¹, C(O)R¹¹, C(=O)NR¹¹₂, S(O)₂R¹¹, S(O)₂NR¹¹₂, or -CN

20 In yet a preferred embodiment of the present invention, V is C₁-C₆ alkyl, aryl or heteroaryl said group being substituted with 0-3 R¹¹, 0-3 R¹² and 0-3 R¹⁵ or V is C₁-C₃ alkylene-NR¹¹C(O)R¹⁴ or C₁-C₃ alkylene-NR¹¹C(O)OR¹⁴ substituted with 0-3 R¹⁵,

25 In yet a preferred embodiment of the present invention, R¹¹ is H or selected independently among the group consisting of C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl, or heteroaryl, said group being substituted with 0-3 R¹² and 0-3 R¹⁵.

30 The function of the carrier is to ensure a high reactivity of the functional entity precursor towards a broad range of carbonyl recipient reactive groups. Substituents on the carrier alter the reactivity of the functional group precursor and can be designed to direct the stereochemically outcome of the reaction by an individual skilled in the art by evaluation of initial attempts. The transferability and the stereoselectivity may be adjusted in response to the chemical composition of the functional entity precursor, to the nature of the complementing element, to the conditions under which the transfer and recognition is performed, etc.

In a preferred embodiment, the carrier is:




wherein R² is a valence bond or arylene.

5 The S-C-connecting group provide a means for connecting the Spacer and the Carrier. As such it is primarily of synthetic convenience and does not influence the function of a building block. In a preferred embodiment of the present invention, the S-C connecting group is -S-S-, C₁-C₆ alkylene-S-S-, -C(=O)-NH-(C₁-C₆ alkylene)-, -C(=O)-, or -C(=O)-Arylene-C(R¹⁵)₂-NR¹⁶-C(=O)-


10

The spacer serves to distance the functional entity precursor to be transferred from the bulky complementing element. Thus, when present, the identity of the spacer is not crucial for the function of the building block. It may be desired to have a spacer which can be cleaved by light. In this occasion, the spacer is provided with e.g. the


15 group




20 In the event an increased hydrophilicity is desired the spacer may be provided with a polyethylene glycol part of the general formula:



25 In a certain aspect of the invention the Spacer is -A-, a group C₁-C₆ alkylene-A-, C₂-C₆ alkenylene-A-, or C₂-C₆ alkynylene-A- optionally substituted with 1 to 3 hydroxy groups, or




said spacer being connected through A to a moiety selected from



- 5 where A is a valence bond, $-NR^{16}$, $-C(O)NR^{16}$, $-NR^{16}-C(O)-$, $-O-$, $-S-$, $-C(O)-O-$ or $-OP(=O)(O)-$; B is a valence bond, $-O-$, $-S-$, $-NR^{16}$, $-C(O)-$ or $-C(O)NR^{16}$, and connects to S-C-connecting group; and n and m independently are integers ranging from 1 to 10.

- 10 In a preferred embodiment the Spacer is a valence bond, C_1-C_6 alkylene-A-, C_7-C_8 alkenylene-A-, C_7-C_8 alkynylene-A-, or




said spacer optionally being connected through A to a moiety selected from




- 15 where A is a valence bond, $-C(O)NR^{16}$, $-NR^{16}$, $-O-$, $-S-$, or $-C(O)-O-$; B is a valence bond, $-O-$, $-S-$, $-NR^{16}$ or $-C(O)NR^{16}$ and connects to S-C-connecting group; R^{16} is selected independently from H, C_1-C_6 alkyl, C_3-C_7 cycloalkyl, aryl or C_1-C_6 alkenylene-aryl and n and m independently are integers ranging from 1 to 10.

- 20 In a more preferred aspect of the invention the spacer is -A-, C_1-C_6 alkylene-A-, C_7-C_8 alkenylene-A-



said spacer being connected through A to a moiety selected from




where A is a valence bond, $-NR^{16}$, $-C(O)NR^{16}$, $-NR^{16}-C(O)-$, $-O-$, $-S-$, $-C(O)-O-$ or $-OP(=O)(O)-$; B is a valence bond, $-S-$, $-NR^{16}$, or $-C(O)-$ and connects to S-C-connecting group;

n and m independently are integers ranging from 1 to 10 and

- 5 R^{16} is selected independently from H, C_1-C_6 alkyl, C_3-C_7 cycloalkyl, aryl or C_1-C_6 alkenylene-aryl, wherein G is H or

Still more preferred the spacer is C_2-C_6 alkenylene-A-,

said spacer being connected through A to a moiety selected from



10 where A is a valence bond, $-C(O)NR^{16}$, $-NR^{16}-C(O)-$, $-S-$, $-C(O)-O-$ or $-OP(=O)(O)-$; B is a valence bond, $-S-$, $-NR^{16}$, or $-C(O)-$ and connects to S-C-connecting group;


n and m independently are integers ranging from 1 to 10 and

- 15 R^{16} is selected independently from H, C_1-C_6 alkyl, C_3-C_7 cycloalkyl, aryl or C_1-C_6 alkenylene-aryl; and the spacer is connected to the complementing element through a nucleobase.


The spacer may be attached to the complementing element in any appropriate way.




- 20 Though it is preferred that the spacer is attached to the 5-position of a pyrimidine type nucleobase or the 7-position of a purine or 7-deaza-purine type nucleobase.

In aspect of the invention it is preferred that the complementing element is connected to the spacer via a phosphorus group. It is then preferred that spacer is -A-,




said spacer being connected through A to a moiety selected from





5 $\text{-NH-C(=O)-Arylene-C(R}^{16}\text{)}_2\text{-NH-C(=O)-, -C(=O)-, -C(=O)-C}_1\text{-C}_6\text{ alkylene- or -C(=O)-Arylene-C(R}^{16}\text{)}_2\text{-NR}^{18}\text{-C(=O)-, where the right hand side of the formulae connects to the carrier.}$

In another aspect, the **S-C-connecting group** is a valence bond, $\text{-NH-C(=O)-, -NH-C(=O)-C}_1\text{-C}_6\text{ alkylene-, -S-S-, -S-S-C}_1\text{-C}_6\text{ alkylene-, -C(=O)-NH-(C}_1\text{-C}_6\text{ alkylene)-,}$



10 $\text{-NH-C(=O)-Arylene-C(R}^{16}\text{)}_2\text{-NH-C(=O)-, where the right hand side of the formulae connects to the carrier.}$

15 In a preferred aspect of the invention the **S-C-connecting group** is $\text{-S-S-, -C}_1\text{-C}_6\text{ alkylene-S-S-, -C(=O)-NH-(C}_1\text{-C}_6\text{ alkylene)-, -C(=O)-, or -C(=O)-Arylene-C(R}^{16}\text{)}_2\text{-NR}^{18}\text{-C(=O)-, where the right hand side of the formulae connects to the carrier.}$

In a still more preferred aspect of the invention the **S-C-connecting group** is $\text{-S-S-, -C(=O)-, or -C(=O)-Arylene-C(R}^{16}\text{)}_2\text{-NR}^{18}\text{-C(=O)-, where the right hand side of the formulae connects to the carrier.}$

20 The building blocks of the present invention can be used in a method for transferring a functional entity precursor to a recipient reactive group, said method comprising the steps of

providing one or more building blocks as described above and

contacting the one or more building blocks with a corresponding encoding element associated with a recipient reactive group under conditions which allow for a recognition between the one or more complementing elements and the encoding elements, said contacting being performed prior to, simultaneously with, or subsequent to a transfer of the functional entity precursor to the recipient reactive group.

5 The encoding element may comprise one, two, three or more codons, i.e. sequences that may be specifically recognised by a complementing element. Each of the codons may be separated by a suitable spacer group. Preferably, all or at least a majority of the codons of the template are arranged in sequence and each of the codons are separated from a neighbouring codon by a spacer group. Generally, it is preferred to have more than two codons on the template to allow for the synthesis of more complex encoded molecules. In a preferred aspect of the invention the number of codons of the encoding element is 2 to 100. Still more preferred are encoding elements comprising 3 to 10 codons. In another aspect, a codon comprises 1 to 50 nucleotides and the complementing element comprises a sequence of nucleotides complementary to one or more of the encoding sequences.

20 The recipient reactive group may be associated with the encoding element in any appropriate way. Thus, the reactive group may be associated covalently or non-covalently to the encoding element. In one embodiment the recipient reactive group is linked covalently to the encoding element through a suitable linker which may be separately cleavable to release the reaction product. In another embodiment, the reactive group is coupled to a complementing element, which is capable of recognising a sequence of nucleotides on the encoding element, whereby the recipient reactive group becomes attached to the encoding element by hybridisation. Also, the recipient reactive group may be part of a chemical scaffold, i.e. a chemical entity having one or more reactive groups available for receiving a functional entity precursor from a building block.

30 The recipient reactive group may be any group able to cleave the Carrier-Functional Entity Precursor bond to release the functional entity precursor. The recipient reactive group is electrophilic, such as an aldehyde or a ketone. The electrophile usually reacts with an anion formed on the functional entity precursor. The outcome of the

reaction is a transformation of the C=O) recipient reactive group into a C=(Functional entity precursor) moiety.

According to a preferred aspect of the invention the building blocks are used for the formation of a library of compounds. The complementing element of the building block is used to identify the functional entity. Due to the enhanced proximity between reactive groups when the complementing entity and the encoding element are contacted, the functional entity precursor together with the identity programmed in the complementing element is transferred to the encoding element associated with recipient reactive group. Thus, it is preferred that the sequence of the complementing element is unique in the sense that the same sequence is not used for another functional entity. The unique identification of the functional entity enable the possibility of decoding the encoding element in order to determine the synthetic history of the molecule formed. In the event two or more functional entities have been transferred to a scaffold, not only the identity of the transferred functional entities can be determined. Also the sequence of reaction and the type of reaction involved can be determined by decoding the encoding element. Thus, according to a preferred embodiment of the invention, each different member of a library comprises a complementing element having a unique sequence of nucleotides, which identifies the functional entity.

Brief description of the drawings

Figure 1. Two setups for Functional Entity Precursor Transfer

Figure 2. Examples of specific base pairing

Figure 3. Example of non-specific base-pairing

Figure 4. Backbone examples

Figure 5 Three examples of building blocks

Detailed Description of the Invention

A building block of the present invention is characterized by its ability to transfer its functional entity precursor to a recipient reactive group. This is done by transforming the C=O double bond of a scaffold to a C=(Functional entity precursor) moiety.

Two setups for generalized functional entity precursor transfer from a building block are depicted in figure 1. In the first example, one complementing element of a build-

ing block recognizes a coding element carrying another functional entity precursor, hence bringing the functional entities in close proximity. This results in a reaction between functional entity precursor 1 and 2 forming a covalent bond between these concurrent with the cleavage of the bond between functional entity precursor 2 and its carrier. In the second example, a coding element brings together two building blocks resulting in functional entity precursor transfer from one building block to the other.

In library generation of chemical entities the tolerance towards a broad range of building blocks is of utmost importance. Useful building blocks for generation of libraries used for drug discovery should therefore possess the necessary reactivity to enable the transfer of the functional entity precursor to a broad range of reaction counter parts, resulting in generation of a highly diverse library, and not only a limited number of products. The Wittig reaction is a powerful C=C bond forming reaction, but the number of useful building blocks are limited to primarily aldehydes. Therefore, the present invention particularly relates to demonstrate the broadness of building blocks capable of acting as C=C bond forming reagents in a modified Wittig reaction (HWE reaction), which allow reaction with aldehydes as well as ketones. Furthermore, use of HWE-building blocks opens up the possibility of directing the stereochemically outcome of the reaction by carefully selecting the substituents on the carrier of the building block.

Figure 5 illustrates three specific compounds according to the invention. For illustrative purposes the individual features used in the claims are indicated. The upper compound is an example of a building block wherein the linker is backbone attached at the 3'-position. The first part of the linker, i.e. the spacer, is an aliphatic chain ending in a nitrogen atom. The nitrogen atom bridges to the S-C-connecting group, which is an N-acylated arylmethyleamine. The carrier attached to the left hand side carbonyl group of the S-C-connecting group is a six-membered phosphinane structure. The phosphor atom is attached to the functional entity precursor, which is an aliphatic ester. When the building block is presented to a recipient reactive group, such as aldehyde or a ketone, the aliphatic ester is transferred forming the C=C double bond of an $\alpha - \beta$ unsaturated ester.

The middle compound illustrates a 5' attachment of a linker. The linker is linked through a phosphate group and extends into a three membered aliphatic chain.

Through another phosphate group and a PEG linker the complementing element is linked via an amide bond to the Carrier. When the building block is presented to a ketone or an aldehyde the functional entity precursor is transferred forming an $\alpha - \beta$ unsaturated ester.

5

The lower compound illustrates a nucleobase attachment of the linker. The linker attaches to the 5' position of a pyrimidine type nucleobase and extends through an $\alpha - \beta$ unsaturated N-methylated amide to the S-C-connecting group, which is a 4-amino methyl benzoic acid derivative. The functional entity precursor can be transferred to a recipient reactive group forming an $\alpha - \beta$ unsaturated keton carrying a protected amine group.


10

Assembly of building blocks

The Carrier-Functional Entity Precursor ensemble may be bound to the Spacer by several different reactions.

15

Formation of an amide bond between a carboxylic acid of the Carrier and an amine group of a Spacer



Examples of Carrier-Functional entity reagents:

	<p>$X = \text{NR}, \text{O}, \text{S}, \text{Se};$ where each of the individual X is chosen independently</p> <p>$R = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl}, \text{aryl or heteroaryl}$</p> <p>$R_1 = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_1\text{-C}_6 \text{ hydroxyalkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl or aryl};$ where each of the R_1 groups are chosen independently</p> <p>$R_2 = \text{a valence bond}, \text{C}_1\text{-C}_6 \text{ Alkylene}, \text{C}_2\text{-C}_6 \text{ Alkenylene}, \text{C}_2\text{-C}_6 \text{ Alkynylene}, \text{Arylene}, \text{Heteroarylene}$</p>
--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

SUBSTITUTE SHEET (RULE 26)

	<p>roarylene, $\text{C}_1\text{-C}_6 \text{ Alkylene-arylene or } \text{C}_1\text{-C}_6 \text{ Alkylene-heteroarylene}$</p> <p>$X = \text{NR}, \text{O}, \text{S}, \text{Se};$ where each of the individual X is chosen independently</p> <p>$R = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl}, \text{aryl or heteroaryl}$</p> <p>$R_1 = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_1\text{-C}_6 \text{ hydroxyalkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl or aryl};$ where each of the R_1 groups are chosen independently</p> <p>$R_2 = \text{a valence bond}, \text{C}_1\text{-C}_6 \text{ Alkylene}, \text{C}_2\text{-C}_6 \text{ Alkenylene}, \text{C}_2\text{-C}_6 \text{ Alkynylene}, \text{Arylene}, \text{Heteroarylene}, \text{C}_1\text{-C}_6 \text{ Alkylene-arylene or } \text{C}_1\text{-C}_6 \text{ Alkylene-heteroarylene}$</p>
	<p>$X = \text{NR}, \text{O}, \text{S}, \text{Se};$ where each of the individual X is chosen independently</p> <p>$R = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_1\text{-C}_6 \text{ alkenyl}, \text{C}_1\text{-C}_6 \text{ alkynyl}, \text{aryl or heteroaryl}$</p> <p>$R_1 = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_1\text{-C}_6 \text{ hydroxyalkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl or aryl};$ where each of the R_1 groups are chosen independently</p> <p>$R_2 = \text{a valence bond}, \text{C}_1\text{-C}_6 \text{ Alkylene}, \text{C}_2\text{-C}_6 \text{ Alkenylene}, \text{C}_2\text{-C}_6 \text{ Alkynylene}, \text{Arylene}, \text{Heteroarylene}, \text{C}_1\text{-C}_6 \text{ Alkylene-arylene or } \text{C}_1\text{-C}_6 \text{ Alkylene-heteroarylene}$</p>
	<p>$X = \text{NR}, \text{O}, \text{S}, \text{Se};$ where each of the individual X is chosen independently</p> <p>$R = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl}, \text{aryl or heteroaryl}$</p> <p>$R_1 = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_1\text{-C}_6 \text{ hydroxyalkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl}, \text{aryl}, \text{Halogen}, \text{NO}_2, \text{CN}, \text{C}(\text{Halogen})_3, \text{C}(\text{OR})^n, \text{C}(\text{ONHR})^n, \text{C}(\text{ONR})^n, \text{NC}(\text{OR})^n, \text{S}(\text{O})_2\text{NHR}^n, \text{S}(\text{O})_2\text{NR}^n, \text{S}(\text{O})_2\text{R}^n, \text{P}(\text{O})_2\text{R}^n, \text{P}(\text{O})\text{R}^n, \text{S}(\text{O})\text{R}^n, \text{P}(\text{O})\text{OR}^n, \text{S}(\text{O})\text{OR}^n, \text{N}^+\text{R}^n$</p> <p>$R_2 = \text{a valence bond}, \text{C}_1\text{-C}_6 \text{ Alkylene}, \text{C}_1\text{-C}_6 \text{ Alkylene-heteroarylene}$</p>

SUBSTITUTE SHEET (RULE 26)

$ \begin{array}{c} R^1-X-O \\ \\ R^2-X-P \\ \\ COOH \end{array} $	Alkenylene, C ₁ -C ₆ Alkynylene, Arylene, Heteroarylene, C ₁ -C ₆ Alkylene-arylene or C ₁ -C ₆ Alkylene-heteroarylene
$ \begin{array}{c} R^1-X-O \\ \\ R^2-X-P \\ \\ COOH \end{array} $	X = NR, O, S, Se; where each of the individual X is chosen independently R = H, C ₁ -C ₆ alkyl, C ₁ -C ₆ alkenyl, C ₁ -C ₆ alkynyl, aryl or heteroaryl R ¹ = H, C ₁ -C ₆ alkyl, C ₁ -C ₆ hydroxyalkyl, C ₂ -C ₆ alkenyl, C ₂ -C ₆ alkynyl, aryl, heteroaryl, or a combination thereof R ² = a valence bond, C ₁ -C ₆ Alkylene, C ₁ -C ₆ Alkenylene, C ₁ -C ₆ Alkynylene, C ₁ -C ₆ Alkenylene, Arylene, Heteroarylene, C ₁ -C ₃ Alkylene substituted with 1-7 halogens selected from the group consisting of -F, -Cl or -Br, C ₁ -C ₆ Alkylene-arylene or C ₁ -C ₆ Alkylene-heteroarylene

According to the invention, the functional entity precursor is of the formula C(H)(V)-

W. In an aspect of the invention V is independently H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₆ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of Sn(R⁵)₂R⁷, Sn(OR⁵)₂R⁷, Sn(OR⁵)(OR⁶)R⁷, Br(R⁵)₂R⁷, B(OR⁵)₂R⁷, B(OR⁵)(OR⁶)R⁷, halogen, CN, CNO, C(halogen)₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, N₃, NR⁵R⁶, N⁺R⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, NC, P(=O)(OR⁵)OR⁶, P⁺R⁵R⁶R⁷, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵NR⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶, W is selected among the group consisting of H, aryl, heteroaryl, C(=O)OR⁵, C(OR⁵)₂, C(=O)NR⁵, C(=O)NR⁵OR⁶, C(=NR⁵)OR⁶, C(=NOR⁵)R⁶, C(=NNR⁵)R⁶, S(OR⁵)₂, S(OR⁵)R⁶, S(O)NR⁵, S(O)₂NR⁵, -CN, P(OR⁵)₂, -NO₂, NR⁵⁺ or SR⁵⁺ wherein,

R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₆ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may

together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring.

In another preferred embodiment,

5 V independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, C(halogen)₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, P(=O)(OR⁵)OR⁶, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶, C(=O)NR⁵NR⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶,

10 wherein, R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring.

15 In still another preferred embodiment, V independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, P(=O)(OR⁵)OR⁶, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶, C(=O)NR⁵NR⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶, wherein,

20 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, P(=O)(OR⁵)OR⁶, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶, C(=O)NR⁵NR⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶, wherein,

25 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, P(=O)(OR⁵)OR⁶, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵R⁶), C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶, C(=O)NR⁵NR⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶, wherein,

30 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring.

In still another preferred embodiment,

V independently is H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

5 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring,

10 In still another preferred embodiment,
V independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl, morpholinyl, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

20 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

25 In still another preferred embodiment,
V independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

30 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 mem-

bered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

5 V independently is H, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

10 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

15 In still another preferred embodiment,

V independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

20 R⁵, R⁶, R⁷ and R⁸ independently is H, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

25 V independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R^5 , R^6 , R^7 and R^8 independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

5

In still another preferred embodiment,

V independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

10

R^5 , R^6 , R^7 and R^8 independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

15

In still another preferred embodiment,

V independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

20

R^5 , R^6 , R^7 and R^8 independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

25

30

In still another preferred embodiment,

V independently is H, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶,

S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R^5 , R^6 , R^7 and R^8 independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

5

In still another preferred embodiment,

V independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

15

R^5 , R^6 , R^7 and R^8 independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

20

In still another preferred embodiment,

V independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

25

R^5 , R^6 , R^7 and R^8 independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

30

35

In still another preferred embodiment,

V independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R⁵, R⁶, R⁷ and R⁸ independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

V independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R⁵, R⁶, R⁷ and R⁸ independently is H, methyl, ethyl, propyl or butyl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

V independently is H, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R⁵, R⁶, R⁷ and R⁸ independently is H, methyl, ethyl, propyl or butyl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together

gether form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

V independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R⁵, R⁶, R⁷ and R⁸ independently is H, methyl, ethyl, propyl or butyl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

V independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R⁵, R⁶, R⁷ and R⁸ independently is H, methyl, ethyl, propyl or butyl and wherein R⁵ and R⁶ may together form a 3-8 membered heterocyclic ring or R⁵ and R⁷ may together form a 3-8 membered heterocyclic ring or R⁶ and R⁷ may together form a 3-8 membered heterocyclic ring,

In still another preferred embodiment,

V independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=O)NR⁵R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁶, wherein,

R^5 , R^6 , R^7 and R^8 independently is H, methyl, ethyl, propyl or butyl and wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring.

5

In still another preferred embodiment,

V independently is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂R⁶, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

10

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

15

In still another preferred embodiment,

V independently is aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

20

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

25

In still another preferred embodiment,

V independently is phenyl, naphthyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

30

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

35

In still another preferred embodiment,

V independently is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂R⁶, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

5

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

10

In still another preferred embodiment,

V independently is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂R⁶, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

15

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

20

In still another preferred embodiment,

V independently is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

25

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl,

30

In still another preferred embodiment,

V independently is aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , =O, =NOR⁵, =NNR⁵R⁶, OR⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, NR⁵R⁶, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, C(=O)R⁵, C(=NOR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶ or R⁵,

NO_2 , NR^5R^6 , $\text{NR}^5\text{C}(=\text{O})\text{OR}^6$, $\text{NR}^5\text{C}(=\text{O})\text{OR}^6$, $\text{NR}^5\text{C}(=\text{O})\text{NR}^6\text{R}^7$, $\text{C}(=\text{O})\text{R}^6$, $\text{C}(=\text{NOR}^5)\text{R}^6$, $\text{C}(=\text{O})\text{OR}^6$, $\text{C}(=\text{O})\text{NR}^5\text{R}^6$, $\text{C}(=\text{O})\text{NR}^5\text{OR}^6$ or R^6 ,

wherein,

R^5 , R^6 , R^7 and R^8 independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolyl or isoquinolyl,

5

In still another preferred embodiment,

V independently is phenyl, naphthyl, thienyl, furyl, pyridyl, quinolyl or isoquinolyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , $=\text{O}$, $=\text{NOR}^5$, $=\text{NNR}^5\text{R}^6$, OR^5 , $\text{S}(=\text{O})\text{R}^5$, $\text{S}(=\text{O})_2\text{R}^5$, $\text{S}(=\text{O})_2\text{NR}^5\text{R}^6$, NO_2 , NR^5R^6 , $\text{NR}^5\text{C}(=\text{O})\text{R}^6$, $\text{NR}^5\text{C}(=\text{O})\text{OR}^6$, $\text{NR}^5\text{C}(=\text{O})\text{NR}^6\text{R}^7$, $\text{C}(=\text{O})\text{R}^5$, $\text{C}(=\text{NOR}^5)\text{R}^6$, $\text{C}(=\text{O})\text{OR}^5$, $\text{C}(=\text{O})\text{NR}^5\text{R}^6$, $\text{C}(=\text{O})\text{NR}^5\text{OR}^6$ or R^6 , wherein,

R^5 , R^6 , R^7 and R^8 independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolyl or isoquinolyl,

15

In still another preferred embodiment,

V independently is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , $=\text{O}$, $=\text{NOR}^5$, $=\text{NNR}^5\text{R}^6$, OR^5 , $\text{S}(=\text{O})\text{R}^5$, $\text{S}(=\text{O})_2\text{R}^5$, $\text{S}(=\text{O})_2\text{NR}^5\text{R}^6$, NO_2 , NR^5R^6 , $\text{NR}^5\text{C}(=\text{O})\text{R}^6$, $\text{NR}^5\text{C}(=\text{O})\text{OR}^6$, $\text{NR}^5\text{C}(=\text{O})\text{NR}^6\text{R}^7$, $\text{C}(=\text{O})\text{R}^5$, $\text{C}(=\text{NOR}^5)\text{R}^6$, $\text{C}(=\text{O})\text{OR}^5$, $\text{C}(=\text{O})\text{NR}^5\text{R}^6$, $\text{C}(=\text{O})\text{NR}^5\text{OR}^6$ or R^6 , wherein,

20

R^5 , R^6 , R^7 and R^8 independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolyl or isoquinolyl,

25

In still another preferred embodiment,

V independently is thienyl, furyl, pyridyl, quinolyl or isoquinolyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , $=\text{O}$, $=\text{NOR}^5$, $=\text{NNR}^5\text{R}^6$, OR^5 , $\text{S}(=\text{O})\text{R}^5$, $\text{S}(=\text{O})_2\text{R}^5$, $\text{S}(=\text{O})_2\text{NR}^5\text{R}^6$, NO_2 , NR^5R^6 , $\text{NR}^5\text{C}(=\text{O})\text{R}^6$, $\text{NR}^5\text{C}(=\text{O})\text{OR}^6$, $\text{NR}^5\text{C}(=\text{O})\text{NR}^6\text{R}^7$, $\text{C}(=\text{O})\text{R}^5$, $\text{C}(=\text{NOR}^5)\text{R}^6$, $\text{C}(=\text{O})\text{OR}^5$, $\text{C}(=\text{O})\text{NR}^5\text{R}^6$, $\text{C}(=\text{O})\text{NR}^5\text{OR}^6$ or R^6 , wherein,

30

R^5 , R^6 , R^7 and R^8 independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolyl or isoquinolyl,

35

In still another preferred embodiment,

V independently is H, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cycloheteroalkyl, aryl or heteroaryl

5

In still another preferred embodiment,

V independently is H,

In still another preferred embodiment,

V independently is $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl or $\text{C}_3\text{-C}_7$ cycloheteroalkyl,

10

In still another preferred embodiment,

V independently is methyl, ethyl, propyl or butyl

In still another preferred embodiment

V independently is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl

15

In still another preferred embodiment

V independently is aziridinyl, pyrrolidinyl, piperidinyl or morpholinyl

20

In still another preferred embodiment,

V independently is aryl or heteroaryl

In still another preferred embodiment,

V independently is phenyl or naphthyl

25

In still another preferred embodiment,

V independently is thienyl, furyl, pyridyl, quinolyl or isoquinolyl

In yet another preferred embodiment,

W is selected among the group consisting of $\text{C}(=\text{O})\text{OR}^5$, $\text{C}(=\text{O})\text{R}^5$, $\text{C}(=\text{O})\text{NR}^5$, $\text{C}(=\text{O})\text{NR}^5\text{OR}^6$, $\text{C}(=\text{NR}^5)\text{OR}^6$, $\text{C}(=\text{NOR}^5)\text{R}^6$, $\text{C}(=\text{NNR}^5_2)\text{R}^6$, $\text{S}(=\text{O})_2\text{R}^5$, $\text{S}(=\text{O})_2\text{NR}^5_2$, and $-\text{CN}$

30


In yet another preferred embodiment,

35

W is selected among the group consisting of C(=O)OR⁵, C(O)R⁵, C(=O)NR⁵, S(O)₂R⁵, S(O)₂NR⁵, and -CN

Experimental section

Example 1. Synthesis of an aliphatic carboxylic acid Carrier Precursor (I):



5

2,2-Bis(hydroxymethyl)-propionic acid (13.41 g, 0.01 mol), acetone (200 ml, 2.72 mol), molecular sieves (10 g) and sulphuric acid (0.5 ml, catalytic amounts) were mixed in the given order at rt. The suspension was refluxed for 10 hours, then neutralized with sodium hydrogen carbonate and filtered through activated carbon on a plough of silica. Evaporation furnished 2,2,5-trimethyl-5-carboxy-1,3-dioxan as a crystalline solid 15.23 g (87%). ¹H-NMR (MeOH-d₄): δ 4.17 (d, 2H), 3.68 (d, 2H), 1.42 (s, 3H), 1.35 (s, 3H), 1.16 (s, 3H).

10

2,2,5-trimethyl-5-carboxy-1,3-dioxan (15.23 g, 87.4 mmol) was dissolved in THF (60 ml) and added DCC (dicyclohexyl carbodiimide, 18.04 g, 87.4 mmol), followed by slow addition of benzyl alcohol (9.04 ml, 87.4 mmol) at 0 °C. Stirring was continued at rt. for 18 hours. The mixture was filtered and solvent was removed by evaporation in vacuo. Distillation followed by chromatography (Ethyl acetate Heptane 1:3) gave 4.1 g of 2,2,5-trimethyl-5-carboxy-1,3-dioxan benzyl ester as crystals. ¹H-NMR (CDCl₃): δ 7.36 (m, 5H), 5.20 (s, 2H) 4.24 (d, 2H), 3.67 (d, 2H), 1.47 (s, 3H), 1.40 (s, 3H), 1.22 (s, 3H).

15

20


2,2,5-Trimethyl-5-carboxy-1,3-dioxan benzyl ester (4.02 g, 15.2 mmol) was dissolved in a mixture of acetone: water : acetic acid (20 ml : 8 ml : 2 ml). The solution was heated at 45 °C for 10 hours and at 56 °C for additional 10 hours. The temperature was raised to 80 °C and the acetone was slowly replaced with water. Stirring was continued for 1.5 hour at 80 °C. Concentration twice from toluene gave 2,2-bis(hydroxymethyl)-propionic acid benzyl ester (I) as colourless crystals 3.40 g. ¹H-NMR (MeOH-d₄): δ 7.36 (m, 5H), 5.17 (s, 2H), 3.73 (d, 2H), 3.67 (d, 2H), 1.20 (s, 3H).

25

Example 2. Synthesis of a benzoic acid Carrier Precursor (II):


30

34



Isopropylidene-2,2-bis(hydroxymethyl)propionic acid:


5



2,2-Bis(hydroxymethyl)propionic acid (0.12 mol, 15.9 g) was refluxed in acetone (250 mL) with molecular sieves and conc. sulphuric acid (0.5 mL) for 10 hours. The reaction mixture was then neutralised with NaHCO_3 (1M aq.), stirred with activated charcoal and filtered. The product was collected as a white crystalline upon concentration of the solvent.

Yield 50 % (10.5g): $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): 1.07 (s, 3H, $-\text{CH}_3$); 1.26 (s, 3H, $-\text{CH}_3$); 1.34 (s, 3H, $-\text{CH}_3$); 3.53 and 3.57 (d, 2H, $-\text{CH}_2-$); 3.99 and 4.02 (d, 2H, $-\text{CH}_2-$).

15 4-(Boc-amino-methyl)-benzoic acid:




4-Methylaminobenzoic acid was dissolved in dioxane (10 mL) and NaOH (22 mL, 1M solution) and cooled to 0 °C. Di-tert-butyl dicarbonate (10 mmol, 2.18 g) and

35

NaOH (8 mL, 2M solution) was added, and the reaction mixture was left over night at room temperature. Half of the solvent was removed under reduced pressure and ethylacetate added (25 mL). The reaction mixture was then neutralised by adding HCl (2 M solution) to pH = 4, and extracted with ethyl acetate (3×75 mL). The organic phase was dried, and evaporated to dryness, and the product was obtained as a white crystalline solid.

Yield: 65 % (6.0 mmol, 1.51 g): $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): 12.84 (s, 1H); 7.89 (d, 2H); 7.46 (t, 1H); 7.34 (d, 2H); 4.19 (d, 2H); 1.40 (s, 9H).

10 4-[(Boc-amino)-methyl]-benzoic acid benzyl ester:




4-[(Boc-amino)-methyl]-benzoic acid (5.89 mmol, 1.48 g) in anhydrous DMF (20 mL) was added Cs_2CO_3 (2.95 mmol, 0.96 g) and stirred for 1h at room temperature.

Benzyl bromide (8.2 mmol, 1.0 mL) was added, and the reaction stirred for 9 hours. The solvent was removed under reduced pressure, and the crude was suspended in water (100 mL) and extracted with diethyl ether (3×100 mL). The organic phase was then dried, evaporated to dryness and the obtained product was purified using dry column vacuum chromatography.

Yield = 81 % (4.79 mmol, 1.56 g): $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): 7.95 (d, 2H); 7.48-7.37 (m, 7H); 5.35 (s, 2H); 4.20 (d, 2H); 1.39 (s, 9H).

20 4-Methylamino benzoic acid benzyl ester:




N-Boc-4-methylamino benzoic acid benzyl ester (4.79 mmol, 1.55 g) was dissolved in DCM (25 mL) with TFA (10 % v/v) and triethylsilane (1 % v/v) and stirred for 30 minutes. The solvent was removed under reduced pressure and the product purified using dry column vacuum chromatography.

30

36

Yield = 47 % (2.28 mmol, 550 mg); ¹H-NMR (DMSO-d₆): 8.69 (s, 2H); 8.03 (d, 2H); 7.62 (d, 2H); 7.50-7.36 (m, 5H); 5.37 (s, 2H); 4.14 (s, 2H).


5 4-[(2,2,5-Trimethyl-1,3)dioxane-5-carbonyl]-amino]-methyl)-benzoic acid benzyl ester



10 Isopropylidene-2,2-bis(hydroxymethyl)propionic acid (4.10 mmol, 714 mg) and 4-methylamino benzyloxy benzoic acid (4.14 mmol, 1.0 g) in DCM (20 mL) was cooled to 0 °C and diisopropyl carbodiimide (5.5 mmol, 0.7 mL) was added. The reaction mixture was left over night at room temperature, and the solvent was removed under reduced pressure. The crude was dissolved in toluene and filtered. The filtrate was purified using Dry Column Vacuum Chromatography.

15 Yield = 29 % (478 mg); ¹H-NMR (DMSO-d₆): 8.25 (t, 1H); 7.93 (d, 2H); 7.47-7.35 (m, 9H); 5.34 (s, 2H); 4.39 (d, 2H); 4.04 (d, 2H); 3.65 (d, 2H); 1.37 (s, 3H); 1.29 (s, 3H); 1.05 (s, 3H);

4-[(3-Hydroxy-2-hydroxymethyl-2-methyl-propionylamino)-methyl]-benzoic acid benzyl ester




20

4-[(2,2,5-Trimethyl-1,3)dioxane-5-carbonyl]-amino]-methyl)-benzoic acid benzyl ester (1.2 mmol, 478 mg) was dissolved in acetic acid (11.5 mL, 87 % v/v) and stirred at 40 °C for 3 hours. The product II was obtained by evaporation of the reaction mixture under reduced pressure and co evaporation from anhydrous toluene (3×20 mL).

25 Yield = 90 %; ¹H-NMR (DMSO-d₆): 8.07 (t, 1H); 7.92 (d, 2H); 7.48-7.12 (m, 7H); 5.34 (s, 2H); 4.72 (bs, 2H); 4.37 (d, 2H); 3.46 (m, 4H); 1.04 (s, 3H).

30 **Example 3.** Synthesis of an aliphatic carboxylic acid Carrier-Functional entity precursor ensemble (III):

37



III


2,2-Bis(hydroxymethyl) propionic acid benzyl ester (1.01 g, 4.5 mmol) was dissolved in toluene (10 ml) and triethylamine (0.75 ml, 1.2 eq.). Methyl dichlorophospho-

5 phorylacetate (0.55 ml, 0.45 mmol, prepared according to the procedure described in *J. Org. Chem.* (1997), 62, 1934-39) was added slowly at 0 °C. Stirring was continued at 0 °C for 15 min. and at rt. for 60 min. The precipitate was filtered of and the filtrate was diluted with ethylacetate (30 ml) and successively washed with, 1 M citric acid (10 ml), 1M sodium hydroxide (2 x 10 ml), saturated ammonium chloride (2 x 10 ml) and saturated sodium chloride (2 x 10 ml), dried with magnesium sulphate and evaporation gave 1.00 g crude material. Purification on silica EtOAc Heptane 1:1 → EtOAc furnished 300 mg of a diastereomeric mixture of 2-methoxycarbonylmethyl-5-methyl-2-oxo-2i5-[1,3,2]dioxaphosphinane-5-carboxylic acid benzyl ester. A) ¹H-NMR (CDCl₃): δ 7.36 (m, 5H), 5.21 (s, 2H), 4.80 (t, 2H), 4.21 (t, 2H), 3.77 (s, 3H), 3.15 (d, 2H), 1.32 (s, 3H). B) ¹H-NMR (CDCl₃): δ 7.36 (m, 5H), 5.24 (s, 2H), 4.68 (dd, 2H), 4.37 (dd, 2H), 3.73 (s, 3H), 2.99 (d, 2H), 1.20 (s, 3H).

2-Methoxycarbonylmethyl-5-methyl-2-oxo-2i5-[1,3,2]dioxaphosphinane-5-carboxylic acid benzyl ester (100 mg, 0.29 mmol, diastereomer A) was dissolved in methanol (5 ml) and placed inside an autoclave. Air was replaced with argon and the catalyst was added (5% Pd/C, 10 mg). The reaction was stirred in a H₂ atmosphere (50 bar) at rt. for 18 hours. The catalyst was filtered of and the solvent removed to give 71 mg of 2-methoxycarbonylmethyl-5-methyl-2-oxo-2i5-[1,3,2]dioxaphosphinane-5-carboxylic acid (III), as colourless crystals. ¹H-NMR (D₂O): δ 4.74 (d, 2H), 4.35 (dd, 2H), 3.72 (s, 3H), 3.41 (d, 2H), 1.10 (s, 3H)

25

Example 4: Synthesis of a benzoic acid Carrier-Functional entity precursor ensembles(IV):



30 PCl₅ (75 mmol) was added to the phosphonate ester(30.0 mmol) at 0 °C. After stirring for 15 min. the temperature was raised to 75 °C for 10 hours. The reaction was

followed by ³¹P-NMR. Distillation at reduced pressure yielded the corresponding dichloride products in 40-50% yield.


Ethyl 2-(dichlorophosphoryl)propionate (bp at 4 mbar: 123 – 125 °C). MS found m/e

5 219 [M+H]⁺

Ethyl 2-(dichlorophosphoryl)butyrate (bp at 4 mbar: 125 – 127 °C). MS found m/e

233 [M+H]⁺

10



15 4-[(3-Hydroxy-2-hydroxymethyl-2-methyl-propionylamino)-methyl]-benzoic acid benzyl ester (1 mmol) was dissolved in toluene triethylamine (2.5 mmol), heating required. The solution was cooled until a precipitate emerged and solution of alkyl 2-(dichlorophosphoryl)carboxylate (1 mmol) in toluene was added. The mixture was stirred at 0 °C for approximately 2 hours and left at rt over night. Ethylacetate (20 mL) was added and the precipitated triethyl ammonium chloride was filtered off. The filtrate was subsequently washed with citric acid (1M, 10 mL), sodium hydroxide (1M, 2 x 10 mL), saturated ammonium chloride (10 mL), dried with magnesium sulphate and evaporated to an oil which was purified by HPLC 0.1% HCCOH, 99.9% H₂O → 0.1% HCCOH, 99.9% MeCN. Fractions containing the desired mass were collected and evaporated to give the desired products in 40-50% yield

25

4-[(2-(1-Ethoxycarbonyl-ethyl)-5-methyl-2-oxo-2λ⁵-[1,3,2]dioxaphosphinane-5-carboxyl)-amino]-methyl)-benzoic acid benzyl ester. MS m/e 504 [M+1]⁺

30

4-[(2-(1-Ethoxycarbonyl-propyl)-5-methyl-2-oxo-2λ⁵-[1,3,2]dioxaphosphinane-5-carboxyl)-amino]-methyl)-benzoic acid benzyl ester. MS m/e 518 [M+1]⁺

SUBSTITUTE SHEET (RULE 26)

Debenzylation were performed by dissolving the purified benzylesters (0.5 mmol) in methanol (5 mL) and placed inside an autoclave. Air was replaced with argon and the catalyst was added (20% Pd/C, 50 mg). The reaction was stirred in a H₂ atmosphere (60 bar) at 40 °C for 2 hours. The catalyst was filtered off and the solvent removed *in vacuo*. The pure products were obtained by HPLC purification using 0.1% HCCOH, 99.9% H₂O → 0.1% HCCOH, 99.9% MeCN. Fractions containing the desired mass were collected and evaporated to dryness.

5


10 **Example 4a**

4-[(2-(1-Ethoxycarbonyl-ethyl)-5-methyl-2-oxo-2λ⁵-[1,3,2]dioxaphosphinane-5-carboxyl)-amino]-methyl)-benzoic acid. MS m/e 414 [M+1]⁺

Example 4b

15 4-[(2-(1-Ethoxycarbonyl-propyl)-5-methyl-2-oxo-2λ⁵-[1,3,2]dioxaphosphinane-5-carboxyl)-amino]-methyl)-benzoic acid. MS m/e 428 [M+1]⁺

20 **General Procedure 1: Preparation of building blocks by loading a Carrier-Functional entity ensemble onto a nucleotide derivative comprising an amino group:**



15 μL of a 150 mM building block solution of FE¹-Carrier-COOH is mixed with 15 μL of a 150 mM solution of EDC and 15 μL of a 150 mM solution of N-hydroxysuccinimide (NHS) using solvents like DMF, DMSO, water, acetonitril, THF, DCM, methanol, ethanol or a mixture thereof. Optionally, the NHS is replaced by a base such as TEA, DIEA, pyridine or DMAP. The mixture is left for 15 min at 25 °C. 45 μL of an aminoaligo (10 nmol) in 100 mM buffer at a pH between 5 and 10, preferably 6.0-7.5 is added and the reaction mixture is left for 2 hours at 25 °C. Excess building block and organic by-products were removed by extraction with EtOAc (400 μL). Remaining EtOAc is evaporated *in vacuo* using a speedvac. The building block is purified following elution through a BioRad micro-spin chromatography column, and analyzed by electron spray mass spectrometry (ESI-MS).

30

SUBSTITUTE SHEET (RULE 26)


Amino-oligo's used:

A: 5'-CTA GGG ACG AGC ATC CAT CG X

X = Amino-C2-dT phosphate (Glen catalogue # 10-1037-)

Example 5 (General Procedure 1)


5 Oligo A with 5' Biotin (Glen catalogue # 10-5950) loaded with Example 3



MS (calc.) = 7246.92; MS (found) = 7246.97

Example 6 (General Procedure 1)


10 Oligo A loaded with example 4a



15 MS (calc.) = 7004.71; MS (found) = 7002.03

Example 7 (General Procedure 1)

Oligo A loaded with example 4b



MS (calc.) = 7018.74; MS (found) = 7016.52

5 Use of building blocks


Loading of a carbonyl containing scaffolds on an oligonucleotide containing an amino group:

25 μ L of a 150 mM carbonyl containing benzoic acid derivative in DMF was mixed with 25 μ L of a 150 mM solution of EDC in DMF. The mixture was left for 30 min at 25°C. 50 μ L of an amino-oligo (10 nmol) in 100 mM HEPES buffer pH 7.5 was added and the reaction mixture was left for 20 min at 25°C. The excess building block was removed by extraction with EtOAc (500 μ L) and remaining EtOAc was removed *in vacuo* by spinning 10 min in a speedvac. The amino-oligo loaded with the benzoic acid derivative was ethanol precipitated twice using NH_4OAc and analysed by electrospray mass spectrometry (ES-MS).

B: 5'-XGATGGATGCTCGTCCCTAG

X = 5'-amine-C6 (Glen catalogue # 10-1906-)


20 Loading of 3-formyl benzoic acid scaffold on oligo B:



MS (calc.) = 6420.32 MS (found) = 6418.59

25


General Procedure 2: Transfer of a functional entity from a building block to a scaffold carrying a reactive recipient group:



5 An oligonucleotide loaded with a phosphonate ester derivative is combined at 10 μ M final concentration with one equivalent of a complementary DNA template displaying an aldehyde or ketone. Olefination proceeds at 37 °C for 12 hours in 50 mM sodium borate, pH 9 buffer. Organic by-products are removed by extraction with EtOAc (400 μ L), followed by evaporation of residual organic solvent for 10 min *in vacuo* using a speedvac. Oligonucleotides are isolated by eluting sample through a BioRad micro-spin chromatography column. Products are characterized by ES-MS analysis.

10


Example 8 (General Procedure 2)



Oligo = Oligo B

15 The compound was obtained reacting example 5 with scaffold oligo V
MS (calc.) = 6476.39 MS (found) = 6474.29

Example 9 (General Procedure 2)




Oligo = Oligo B

Oligo = Oligo B

The compound was obtained reacting example 6 with scaffold oligo V
MS (calc.) = 6504.44 ## MS (found) = 6502.65

Example 10 (General Procedure 2)



5

Oligo = Oligo B

The compound was obtained reacting example 7 with scaffold oligo V
MS (calc.) = 6518.47 MS (found) = 6516.08

10 **Model Example 11.** General route to the formation of Wittig and HWE monomer building blocks and use of these:


Commercially available building block (16) may be transformed into the NHS ester (17) by standard means, *i.e.* DCC or DIC couplings.

15 An amine carrying oligonucleotide in buffer 50 mM MOPS or hepes or phosphate pH 7.5 is treated with a 1-100 mM solution and preferably 7.5 mM solution of the organic building block in DMSO or alternatively DMF, such that the DMSO/DMF concentration is 5-50%, and preferably 10%. The mixture is left for 1-16 h and preferably 2-4 h at 25 °C. To give the phosphine bound monomer building block (18). This monomer building block is further transformed by addition of the appropriate alkyl-
20 halide, *e.g.* *N,N*-dimethyl-2-iodoacetamide as a 1-100 mM and preferably 7.5 mM solution in DMSO or DMF such that the DMSO/DMF concentration is 5-50%, and preferably 10%. The mixture is left for 1-16 h and preferably 2-4 h at 25 °C. To give the monomer building block (19). Alternative to this, may the organic building block (17) be *P*-alkylated with an alkylhalide and then be coupled onto an amine carrying oligonucleotide to yield (19).

25

An aldehyde bound monomer building block (20), *e.g.* formed by the reaction between the NHS ester of 4-formylbenzoic acid and an amine carrying oligonucleotide,

using conditions similar to those described above, will react with (19) under slightly alkaline conditions to yield the alkene (21).



5

The reaction of monomer building blocks (19) and (20) may be conducted as follows:


The template oligonucleotide (1 nmol) is mixed with monomer building block (19) (1 nmol) and (20) (1 nmol) in 0.1 M TAPS, phosphate or hepes-buffer and 1 M NaCl solution, pH=7.5-8.5 and preferably pH=8.0. The reaction mixture is left at 35-65 °C preferably 58 °C over night to yield template bound (21).

As an alternative to (17) phosphonates (24) may be used instead. They may be prepared by the reaction between diethylchlorophosphate (22) and the appropriate car-

10

boxy carrying alcohol. The carboxylic acid is then transformed into the NHS ester (24) and the process and alternatives described above may be applied. Although instead of a simple *P*-alkylation, the phosphite will undergo Arbuzov's reaction and generate the phosphonate. Monomer building block (25) benefits from the fact that it is more reactive than its phosphonium counterpart (19).

5



47

CCC	N,N'-Dicyclohexylcarbodiimide
DIC	Diisopropylcarbodiimide
DIEA	Diisopropylethylamine
TEA	Triethylamine
DNA	Deoxyribonucleic Acid
EDC	1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide HCl
LNA	Locked Nucleic Acid
NHS	N-hydroxysuccinimid
PNA	Peptide Nucleic Acid
RNA	Ribonucleic acid
RP-HPLC	Reverse Phase High Performance Liquid Chromatography
ES-MS	Electron spray mass spectrometry
DMF	Dimethylformamide
THF	Tetrahydrofuran

1. A building block of the general formula


5 **Complementing Element – Linker – Carrier – Functional entity precursor**
capable of transferring a functional entity precursor to a recipient reactive group,
wherein

Complementing Element is a group identifying the functional entity.

Linker is a chemical moiety comprising a spacer and a S-C-connecting

10 group, wherein the spacer is a valence bond or a group distancing the functional entity precursor to be transferred from the complementing element and the S-C-connecting group connects the spacer with the Carrier,

Carrier is selected among the groups consisting of



wherein the Functional entity precursor is attached to the phosphorous atom,

15 each individual X is chosen independently among the group consisting of NR^{16} , O, S, and Se;

$$U = \mathbb{R}^3;$$

$R^1 = H, C_1-C_8 \text{ alkyl}, C_1-C_8 \text{ hydroxyalkyl}, C_2-C_8 \text{ alkenyl}, C_2-C_8 \text{ alkynyl}, \text{aryl or het-}$

20
eroaryl; optionally substituted with one or more F, Cl, Br, I, or CN, and where each of the individual R¹ groups are chosen independently;

R^2 = a valence bond, C_1-C_8 Alkylene, C_2-C_6 Alkenylene, C_2-C_6 Alkynylene, Arylene, Heteroarylene, C_1-C_8 Alkylene-arylene or C_1-C_8 Alkylene-heteroarylene;

R^3 is selected from -H, -OR⁴, -NR⁴, -F, -Cl, -Br, -I, -NO₂, -CN, -C(Halogen)₃,

25 $-\text{C}(\text{O}R^1)-\text{C}(\text{O}NHR^4)-\text{C}(\text{O}NR^4)_2, -\text{NC}(\text{O}R^1)_2, -\text{S}(\text{O})_2NHR^4, -\text{S}(\text{O})_2NR^4, -\text{S}(\text{O})_2R^4, -\text{P}(\text{O})_2R^4, -\text{P}(\text{O})-R^4, -\text{S}(\text{O})-R^4, -\text{S}(\text{O})-OR^1, -\text{S}(\text{O})-OR^4, -N^+R^4$, wherein and each individual R^4 is chosen independently from H, C_1-C_8 alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, or aryl.

R^{16} is selected independently from H, C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, aryl, C_1 - C_8 al-



1-2,3 or 4, 5 or 6, 7 or 8, 9 or 10, 11 or 12, 13 or 14, 15 or 16, 17 or 18, 19 or 20, 21 or 22, 23 or 24, 25 or 26, 27 or 28, 29 or 30, 31 or 32, 33 or 34, 35 or 36, 37 or 38, 39 or 40, 41 or 42, 43 or 44, 45 or 46, 47 or 48, 49 or 50, 51 or 52, 53 or 54, 55 or 56, 57 or 58, 59 or 60, 61 or 62, 63 or 64, 65 or 66, 67 or 68, 69 or 70, 71 or 72, 73 or 74, 75 or 76, 77 or 78, 79 or 80, 81 or 82, 83 or 84, 85 or 86, 87 or 88, 89 or 90, 91 or 92, 93 or 94, 95 or 96, 97 or 98, 99 or 100, 101 or 102, 103 or 104, 105 or 106, 107 or 108, 109 or 110, 111 or 112, 113 or 114, 115 or 116, 117 or 118, 119 or 120, 121 or 122, 123 or 124, 125 or 126, 127 or 128, 129 or 130, 131 or 132, 133 or 134, 135 or 136, 137 or 138, 139 or 140, 141 or 142, 143 or 144, 145 or 146, 147 or 148, 149 or 150, 151 or 152, 153 or 154, 155 or 156, 157 or 158, 159 or 160, 161 or 162, 163 or 164, 165 or 166, 167 or 168, 169 or 170, 171 or 172, 173 or 174, 175 or 176, 177 or 178, 179 or 180, 181 or 182, 183 or 184, 185 or 186, 187 or 188, 189 or 190, 191 or 192, 193 or 194, 195 or 196, 197 or 198, 199 or 200, 201 or 202, 203 or 204, 205 or 206, 207 or 208, 209 or 210, 211 or 212, 213 or 214, 215 or 216, 217 or 218, 219 or 220, 221 or 222, 223 or 224, 225 or 226, 227 or 228, 229 or 230, 231 or 232, 233 or 234, 235 or 236, 237 or 238, 239 or 240, 241 or 242, 243 or 244, 245 or 246, 247 or 248, 249 or 250, 251 or 252, 253 or 254, 255 or 256, 257 or 258, 259 or 260, 261 or 262, 263 or 264, 265 or 266, 267 or 268, 269 or 270, 271 or 272, 273 or 274, 275 or 276, 277 or 278, 279 or 280, 281 or 282, 283 or 284, 285 or 286, 287 or 288, 289 or 290, 291 or 292, 293 or 294, 295 or 296, 297 or 298, 299 or 300, 301 or 302, 303 or 304, 305 or 306, 307 or 308, 309 or 310, 311 or 312, 313 or 314, 315 or 316, 317 or 318, 319 or 320, 321 or 322, 323 or 324, 325 or 326, 327 or 328, 329 or 330, 331 or 332, 333 or 334, 335 or 336, 337 or 338, 339 or 340, 341 or 342, 343 or 344, 345 or 346, 347 or 348, 349 or 350, 351 or 352, 353 or 354, 355 or 356, 357 or 358, 359 or 360, 361 or 362, 363 or 364, 365 or 366, 367 or 368, 369 or 370, 371 or 372, 373 or 374, 375 or 376, 377 or 378, 379 or 380, 381 or 382, 383 or 384, 385 or 386, 387 or 388, 389 or 390, 391 or 392, 393 or 394, 395 or 396, 397 or 398, 399 or 400, 401 or 402, 403 or 404, 405 or 406, 407 or 408, 409 or 410, 411 or 412, 413 or 414, 415 or 416, 417 or 418, 419 or 420, 421 or 422, 423 or 424, 425 or 426, 427 or 428, 429 or 430, 431 or 432, 433 or 434, 435 or 436, 437 or 438, 439 or 440, 441 or 442, 443 or 444, 445 or 446, 447 or 448, 449 or 450, 451 or 452, 453 or 454, 455 or 456, 457 or 458, 459 or 460, 461 or 462, 463 or 464, 465 or 466, 467 or 468, 469 or 470, 471 or 472, 473 or 474, 475 or 476, 477 or 478, 479 or 480, 481 or 482, 483 or 484, 485 or 486, 487 or 488, 489 or 490, 491 or 492, 493 or 494, 495 or 496, 497 or 498, 499 or 500, 501 or 502, 503 or 504, 505 or 506, 507 or 508, 509 or 510, 511 or 512, 513 or 514, 515 or 516, 517 or 518, 519 or 520, 521 or 522, 523 or 524, 525 or 526, 527 or 528, 529 or 530, 531 or 532, 533 or 534, 535 or 536, 537 or 538, 539 or 540, 541 or 542, 543 or 544, 545 or 546, 547 or 548, 549 or 550, 551 or 552, 553 or 554, 555 or 556, 557 or 558, 559 or 560, 561 or 562, 563 or 564, 565 or 566, 567 or 568, 569 or 570, 571 or 572, 573 or 574, 575 or 576, 577 or 578, 579 or 580, 581 or 582, 583 or 584, 585 or 586, 587 or 588, 589 or 590, 591 or 592, 593 or 594, 595 or 596, 597 or 598, 599 or 600, 601 or 602, 603 or 604, 605 or 606, 607 or 608, 609 or 610, 611 or 612, 613 or 614, 615 or 616, 617 or 618, 619 or 620, 621 or 622, 623 or 624, 625 or 626, 627 or 628, 629 or 630, 631 or 632, 633 or 634, 635 or 636, 637 or 638, 639 or 640, 641 or 642, 643 or 644, 645 or 646, 647 or 648, 649 or 650, 651 or 652, 653 or 654, 655 or 656, 657 or 658, 659 or 660, 661 or 662, 663 or 664, 665 or 666, 667 or 668, 669 or 670, 671 or 672, 673 or 674, 675 or 676, 677 or 678, 679 or 680, 681 or 682, 683 or 684, 685 or 686, 687 or 688, 689 or 690, 691 or 692, 693 or 694, 695 or 696, 697 or 698, 699 or 700, 701 or 702, 703 or 704, 705 or 706, 707 or 708, 709 or 710, 711 or 712, 713 or 714, 715 or 716, 717 or 718, 719 or 720, 721 or 722, 723 or 724, 725 or 726, 727 or 728, 729 or 730, 731 or 732, 733 or 734, 735 or 736, 737 or 738, 739 or 740, 741 or 742, 743 or 744, 745 or 746, 747 or 748, 749 or 750, 751 or 752, 753 or 754, 755 or 756, 757 or 758, 759 or 760, 761 or 762, 763 or 764, 765 or 766, 767 or 768, 769 or 770, 771 or 772, 773 or 774, 775 or 776, 777 or 778, 779 or 780, 781 or 782, 783 or 784, 785 or 786, 787 or 788, 789 or 790, 791 or 792, 793 or 794, 795 or 796, 797 or 798, 799 or 800, 801 or 802, 803 or 804, 805 or 806, 807 or 808, 809 or 810, 811 or 812, 813 or 814, 815 or 816, 817 or 818, 819 or 820, 821 or 822, 823 or 824, 825 or 826, 827 or 828, 829 or 830, 831 or 832, 833 or 834, 835 or 836, 837 or 838, 839 or 840, 841 or

30

Functional entity precursor is of the general formula C(H)(V)-W;

wherein V independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloalkenyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of $\text{Sn(R}^5\text{R}^6)_3$, $\text{Sn(OR}^5\text{)R}^6\text{R}^7$, $\text{Sn(OR}^5\text{)(OR}^6\text{)R}^7$, BR^5R^6 , $\text{B(OR}^5\text{)R}^6$, $\text{B(OR}^5\text{)(OR}^6\text{)}$, halogen, CN, CNO, C(halogen)₃, =O, =NOR⁵, =NNR⁵R⁶, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵R⁶, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵R⁶, NO₂, N₃, NR⁵R⁶, N⁺R⁵R⁶R⁷, NR⁵OR⁶, NR⁵NR⁶R⁷, NR⁵C(=O)R⁶, NR⁵C(=O)OR⁶, NR⁵C(=O)NR⁶R⁷, NC, P(=O)(OR⁵)OR⁶, P⁺R⁵R⁶R⁷, C(=O)R⁵, C(=NR⁵)R⁶, C(=NOR⁵)R⁶, C(=NNR⁵)R⁶, C(=O)OR⁵, C(=O)NR⁵R⁶, C(=O)NR⁵OR⁶, C(=O)NR⁵NR⁶R⁷, C(=NR⁵)NR⁶R⁷, C(=NOR⁵)NR⁶R⁷ or R⁶.

10 wherein.

15 R^5 , R^6 and R^7 independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cydoalkyl, cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CNO, C(halogen)₃, =O, =NOR⁵, =NNR⁵, OR⁵, OC(=O)R⁵, OC(=O)OR⁵, OC(=O)NR⁵, SR⁵, S(=O)R⁵, S(=O)₂R⁵, S(=O)₂NR⁵, NO₂, N₃, NR⁵, N⁺R⁵R¹⁰, NR⁵OR⁵, NR⁵NR⁵R⁷, NR⁵C(=O)R⁵, NR⁵C(=O)OR⁵, NR⁵C(=O)NR⁵R¹⁰, NC, P(=O)(OR⁵)OR⁵, P⁺R⁵R⁵R⁷, C(=O)R⁵, C(=NR⁵)R⁵, C(=NOR⁵)R⁵, C(=NNR⁵)R⁵, C(=O)OR⁵, C(=O)NR⁵R⁵, C(=O)NR⁵OR⁵, C(=NR⁵)NR⁵R⁷, C(=NOR⁵)NR⁵R⁷ or C(=O)NR⁵NR⁵R¹⁰, wherein R^5 and R^6 may together form a 3-8 membered heterocyclic ring or R^5 and R^7 may together form a 3-8 membered heterocyclic ring or R^6 and R^7 may together form a 3-8 membered heterocyclic ring,

wherein,

25 R^6 , R^9 and R^{10} independently is H, alkyl, alkenyl, alkynyl, alkydialkyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl and wherein R^6 and R^9 may together form a 3-8 membered heterocyclic ring or R^6 and R^{10} may together form a 3-8 membered heterocyclic ring or R^9 and R^{10} may together form a 3-8 membered heterocyclic ring.

W is selected among the group consisting of H, aryl, heteroaryl, C(=O)OR¹¹, C(O)R¹¹, C(=O)NR¹¹, C(=O)NR¹¹OR¹¹, C(=NR¹¹)OR¹¹, C(=NOR¹¹)R¹¹, C(=NNR¹¹)₂R¹¹, S(O)R¹¹, S(O)₂R¹¹, S(O)NR¹¹, S(O)₂NR¹¹, -CN, P(O)R¹¹, -NO₂, NR¹¹,₃ or SR¹¹,₂.

where each individual R¹¹ is H or selected independently among the group consisting of C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R¹² and 0-3 R¹⁵ and

each individual R¹² is selected independently from -N₃, -CNO, -C(NO₂H)NH₂, -NHOH, -NHR¹NHR¹, -C(OR¹)₄, -SnR¹₃, -B(OR¹)₂, -P(O)(OR¹)₂ or the group consisting of C₂-C₆ alkenyl, C₂-C₈ alkynyl, C₂-C₈ alkydaryl said group being substituted with 0-2 R¹³.

5 where each individual R^{13} is independently selected from $-NO_2$, $-C(O)OR^{14}$, $-C(O)R^{14}$, $-CN$, $-OSiR^{14}_3$, $-OR^{14}$ and $-NR^{14}_2$.

each individual R^{14} , is independently chosen from a group comprising H, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_3-C_7 cycloalkyl, aryl or C_1-C_6 alkylene-aryl substituted with 0-3 substituents independently selected from $-F$, $-Cl$, $-NO_2$, $-R^4$, $-OR^4$, $-SiR^4_3$.

R^{15} is -O-, -F-, -Cl-, Br-, I-, CN-, NO₂-, OR¹⁶, NR¹⁶, C(O)R¹⁶, -NR¹⁶C(O)OR¹⁶, -SR¹⁶, S(O)R¹⁶, -S(O)₂R¹⁶, -COOR¹⁶, -CONR¹⁶, -C(=N)R¹⁶-COOR¹⁷, -SO₂NH- and -NH-SO₂; R¹⁶ is H or alkyl; n = 0, 1, 2.

2. The compound according to claim 1 wherein the Functional entity precursor is of the general formula C(H)(V)-W, in which V is H, -F, -Cl, -Br, -I, -CN, -NO₂ or -OR¹⁴, or selected among the group consisting of C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₁-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, -O-aryl and -O-heteroaryl said group being substituted with 0-3 R¹¹, 0-3 R¹² and 0-3 R¹⁵.

20 or V is C₁-C₃ alkylene-NR¹¹₂, C₁-C₃ alkylene-NR¹¹C(O)R¹⁴, C₁-C₃ alkylene-NR¹¹C(O)R¹⁴, C₁-C₂ alkylene-O-NR¹¹₂, C₁-C₂ alkylene-O-NR¹¹C(O)R¹⁴, C₁-C₂ alkylene-O-NR¹¹C(O)R¹⁴ substituted with 0-3 R¹⁵.

W is selected among the group consisting of H, aryl, heteroaryl, C(=O)OR¹¹, C(O)R¹¹, C(=O)NR¹², C(=O)NR¹¹OR¹¹, C(=NR¹¹)OR¹¹, C(=NOR¹¹)R¹¹.

25 $C(=NNR^{11})_2R^{11}$, $S(O)R^{11}$, $S(O)_2R^{11}$, $S(O)NR^{11}_2$, $S(O)_2NR^{11}_2$, $-CN$, $P(O)R^{11}_2$, $-NO_2$, NR^{11}_3 or SR^{11}_2

where R¹¹ is H or selected independently among the group consisting of C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R¹² and 0-3 R¹⁵ and

30 R¹² is selected independently from -N₃, -CNO, -C(NOH)NH₂, -NHOH, -NHR¹NHR¹, -C(O)R¹, -SnR³₃, -B(OR³)₂, -P(O)(OR³)₂ or the group consisting of C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₁-C₈ alkadienyl said group being substituted with 0-2 R¹³

where R^{13} is independently selected from $-\text{NO}_2$, $-\text{C}(\text{O})\text{OR}^{14}$, $-\text{C}(\text{O})\text{R}^{14}$, $-\text{CN}$, $-\text{OSiR}^{14}_3$, $-\text{OR}^{14}$ and $-\text{NR}^{14}_2$.

- 5 R^{14} is independently chosen from a group comprising H, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_3-C_7 cycloalkyl, aryl or C_1-C_6 alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -NO₂, -R¹, -OR¹, -SiR¹₃, R^{15} is =O, -F, -Cl, -Br, -I, -CN, -NO₂, -OR¹⁴, -NR¹⁴, -NR¹⁴₂, -C(O)R¹⁴, -NR¹⁴₂C(O)OR¹⁴, -SR¹⁴, -S(O)R¹⁴, -S(O)₂R¹⁴, -COOR¹⁴, -C(O)NR¹⁴₂ and -S(O)₂NR¹⁴₂, R^{16} is selected independently from H, C_1-C_6 alkyl, C_3-C_7 cycloalkyl, aryl, C_1-C_6 alky-



kylene-aryl, $-(CH_2)_n-$ or $-(CH_2)_m-$, G is H or C_1-C_6 alkyl and n is 1, 2, 3 or 4.


3. A compound according to claim 2 wherein W is selected among the group consisting of H, aryl, heteroaryl, COOR¹¹, S(O)R¹¹, S(O)₂R¹¹, S(O)NR¹¹₂, S(O)NR¹¹₂, CN, P(O)R¹¹₂, NR¹¹₃⁺ or SR¹¹₂⁺.

4. A compound according to claim 2 wherein W is selected among the group consisting of C(O)OR¹¹, C(O)R¹¹, C(=O)NR¹¹₂, S(O)₂R¹¹, S(O)₂NR¹¹₂, or -CN


5. A compound according to any of claims 2-4 wherein V is C_1-C_6 alkyl, aryl or heteroaryl said group being substituted with 0-3 R¹¹, 0-3 R¹² and 0-3 R¹⁵ or V is C_1-C_6 alkylene-NR¹¹C(O)R¹⁴ or C_1-C_6 alkylene-NR¹¹C(O)OR¹⁴ substituted with 0-3 R¹⁵.

6. A compound according to any of claims 2-5 wherein R¹¹ is H or selected independently among the group consisting of C_1-C_6 alkyl, C_3-C_7 cycloalkyl, aryl, or heteroaryl, said group being substituted with 0-3 R¹² and 0-3 R¹⁵

7. A compound according to claim 1 or 2 wherein Spacer is -A-, a group C_1-C_6 alkylene-A-, C_2-C_6 alkenylene-A-, or C_2-C_6 alkynylene-A- optionally substituted with 1 to 3 hydroxy groups, or




said spacer being connected through A to a moiety selected from



-B-, $-(CH_2)_n-$ -B-, and

8. A compound according to claim 1 or 2 wherein Spacer is a valence bond, C_1-C_6 alky-

alkylene-A-, C_2-C_6 alkenylene-A-, C_2-C_6 alkynylene-A-, or




said spacer optionally being connected through A to a moiety selected from




-B-, $-(CH_2)_n-$ -B-, and

9. A compound according to claim 1 or 2, wherein the spacer is -A-, C_1-C_6 alky-

alkylene-A-, C_2-C_6 alkenylene-A-,





said spacer being connected through A to a moiety selected from




-B-, $-(CH_2)_n-$ -B-, or

10. A compound according to claim 1 or 2, wherein the spacer is -A-, C_1-C_6 alky-


alkylene-A-, C_2-C_6 alkenylene-A-, or C_2-C_6 alkynylene-A-, and

R^{16} is selected independently from H,  or , wherein G is H or C_1-C_8 alkyl.

10. A compound according to claim 1 or 2, wherein the spacer is -A-, C_2-C_8 alkylene-A,





said spacer being connected through A to a moiety selected from




-B-, or $-(CH_2)_n-B-$, where A is a valence bond, $-NR^{16}-C(O)-$, $-O-$, $-S-$, $-C(O)-O-$, or $-OP(=O)(O)-O-$; B is a valence bond, $-S-$, $-NR^{16}$, or $-C(O)-$ and connects to S-C-connecting group;

n and m independently are integers ranging from 1 to 10 and

R^{16} is selected independently from H,  or , wherein G is H or C_1-C_8 alkyl.



11. A compound according to claim 1 or 2, wherein the spacer is C_2-C_8 alkylene-A,

said spacer being connected through A to a moiety selected from




-B-, or $-(CH_2)_n-B-$, where A is a valence bond, $-C(O)NR^{16}$, $-NR^{16}-C(O)-$, $-S-$, $-C(O)-O-$ or $-OP(=O)(O)-O-$; B is a valence bond, $-S-$, $-NR^{16}$, or $-C(O)-$ and connects to S-C-connecting group;

n and m independently are integers ranging from 1 to 10 and


R^{16} is selected independently from H,  or , wherein G is H or C_1-C_8 alkyl; and the spacer is connected to the complementing element through a nucleobase.

12. A compound according to any of the claims 1 to 11, wherein the spacer is attached to the 5-position of a pyrimidine type nucleobase or the 7-position of a purine or 7-deaza-purine type nucleobase.

13. A compound according to claim 1 or 2, wherein the spacer is -A-,





said spacer being connected through A to a moiety selected from





-B-, or $-(CH_2)_n-B-$, where A is a valence bond, $-NR^{16}-C(O)-$, $-O-$, or $-S-$; B is a valence bond, $-S-$, $-NR^{16}$, or $-C(O)-$ and connects to S-C-connecting group;

n and m independently are integers ranging from 1 to 10 and

R^{16} is selected independently from H,  or , wherein G is H or C_1-C_8 alkyl; and the spacer is connected to the complementing element via a phosphorus group.

14. A compound according to claim 13, wherein the phosphorus group is a phosphate or thiophosphate group attached to a 3' or 5' end of a complementing element.


15. A compound according to claims 1 to 14, wherein the S-C-connecting group is a valence bond, $-NH-C(=O)-$, $-NH-C(=O)-C_1-C_8$ alkylene-, $-S-S-$, $-S-S-C_1-C_8$ alkylene-, $-C_1-C_8$ alkylene- $S-S-$, $-C(=O)-NH-(C_1-C_8$ alkylene)-,



$-NH-C(=O)-Arylene-C(R^{16})_2-NH-C(=O)-$, $-C(=O)-$, $-C(=O)-C_1-C_8$ alkylene- or $-C(=O)-$

Arylene-C(R¹⁵)₂-NR¹⁶-C(=O)-, where the right hand side of the formulae connects to the carrier.

16. A compound according to claims 1 to 14, wherein the **S-C-connecting group** is a valence bond, -NH-C(=O)-, -NH-C(=O)-C₁-C₈ alkylene-, -S-S-, -S-S-C₁-C₈ alkylene-, -C(=O)-NH-(C₁-C₈ alkylene)-,




17. A compound according to claims 1 to 14, wherein the **S-C-connecting group** is -NH-C(=O)-Arylene-C(R¹⁶)₂-NH-C(=O)-, where the right hand side of the formulae connects to the carrier.

18. A compound according to claims 1 to 14, wherein the **S-C-connecting group** is -S-S-, -C₁-C₈ alkylene-S-S-, -C(=O)-NH-(C₁-C₈ alkylene)-, -C(=O)-, or -C(=O)-Arylene-C(R¹⁶)₂-NR¹⁶-C(=O)-, where the right hand side of the formulae connects to the carrier.

19. A compound according to claims 1 to 14, wherein the **S-C-connecting group** is -S-S-, -C(=O)-, or -C(=O)-Arylene-C(R¹⁶)₂-NR¹⁶-C(=O)-, where the right hand side of the formulae connects to the carrier.

20. A compound according to claim 1 wherein the carrier is:



wherein R² is a valence bond or arylene.

21. A compound according to claims 1-19 where Complementing element is a nucleic acid.

22. A compound according to claims 1-19 where Complementing element is a sequence of nucleotides selected from the group of DNA, RNA, LNA PNA, morpholino derivatives or combinations thereof.

23. A method for transferring a Functional Entity Precursor to a recipient reactive group, comprising the steps of

providing one or more building blocks according to claims 1 to 21,

contacting the one or more building blocks with a corresponding encoding element associated with a recipient reactive group under conditions which allow for a recognition between the one or more complementing elements and the encoding elements, said contacting being performed prior to, simultaneously with, or subsequent to a transfer of the Functional Entity Precursor to the recipient reactive group.

24. The method according to claim 23, wherein the encoding element comprises one or more encoding sequences comprised of 1 to 50 nucleotides and the one or more complementing elements comprises a sequence of nucleotides complementary to one or more of the encoding sequences.

Figure 1. Two setups for Functional Entity Precursor Transfer




Figure 2. Examples of specific base pairing




Figure 3. Example of non-specific base-pairing

I = Inosine




Figure 4. Backbone examples





Figure 5.



THIS PAGE BLANK (USPT